

# Sevelamer carbonate Rowex 800 mg Film-coated tablets

sevelamer carbonate

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or your pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor. This includes any possible side effects not listed in this leaflet. See section 4.

## What is in this leaflet

1. What Sevelamer carbonate Rowex is and what it is used for
2. What you need to know before you take Sevelamer carbonate Rowex
3. How to take Sevelamer carbonate Rowex
4. Possible side effects
5. How to store Sevelamer carbonate Rowex
6. Contents of the pack and other information

**ROWEX**<sup>®</sup>

## 1. What Sevelamer carbonate Rowex is and what it is used for

Sevelamer carbonate Rowex contains sevelamer carbonate as the active ingredient. It binds phosphate from food in the digestive tract and so reduces serum phosphorus levels in the blood.

Sevelamer carbonate Rowex is used to control hyperphosphataemia (high blood phosphate levels) in:

- adult patients on dialysis (a blood clearance technique). It can be used in patients undergoing haemodialysis (using a blood filtration machine) or peritoneal dialysis (where fluid is pumped into the abdomen and an internal body membrane filters the blood)
- patients with chronic (long-term) kidney disease who are not on dialysis and have a serum (blood) phosphorus level equal to or above 1.78 mmol/l.

Sevelamer carbonate Rowex should be used with other treatments such as calcium supplements and vitamin D to prevent the development of bone disease.

Increased levels of serum phosphorus can lead to hard deposits in your body called calcification. These deposits can stiffen your blood vessels and make it harder for blood to be pumped around the body. Increased serum phosphorus can also lead to itchy skin, red eyes, bone pain and fractures.

## 2. What you need to know before you take Sevelamer carbonate Rowex

**Do not take Sevelamer carbonate Rowex if:**

- you have **low levels of phosphate** in your blood (your doctor will check this for you)
- you have **bowel obstruction**
- you are **allergic to sevelamer carbonate or to any of the other ingredients** of this medicine (listed in section 6).

### Warnings and Precautions

Talk to your doctor before taking Sevelamer carbonate Rowex if any of the following applies to you:

- **swallowing** problems
- problems with **motility (movement) in your stomach and bowel**
- **being sick** frequently
- active **inflammation of the bowel**
- have undergone **major surgery** on your stomach or bowel.

### Additional treatments:

Due to either your kidney condition or your dialysis treatment you may:

- develop low or high levels of calcium in your blood. Since sevelamer carbonate does not contain calcium your doctor might prescribe additional calcium tablets.
- have a low amount of vitamin D in your blood. Therefore, your doctor may monitor the levels of vitamin D in your blood and prescribe additional vitamin D as necessary. If you do not take multivitamin supplements you may also develop low levels of vitamins A, E, K and folic acid in your blood and therefore your doctor may monitor these levels and prescribe supplemental vitamins as necessary.

### Special note for patients on peritoneal dialysis:

You may develop peritonitis (infection of your abdominal fluid) associated

with your peritoneal dialysis. This risk can be reduced by careful adherence to sterile techniques during bag changes. You should **tell your doctor immediately if you experience any new signs or symptoms of abdominal distress, abdominal swelling, abdominal pain, abdominal tenderness, or abdominal rigidity, constipation, fever, chills, nausea or vomiting**. You should expect to be monitored more carefully for problems with low levels of vitamins A, D, E, K and folic acid.

### Children

The safety and efficacy in children (below the age of 6 years) have not been studied. Therefore Sevelamer carbonate Rowex is not recommended for use in children below the age of 6 years.

### Other medicines and Sevelamer carbonate Rowex

Tell your doctor if you are taking or have recently taken or might take any other medicines.

Sevelamer carbonate Rowex should not be taken at the same time as **ciprofloxacin** (an antibiotic).

If you are taking **medicines for heart rhythm problems or for epilepsy**, you should consult your doctor when taking Sevelamer carbonate Rowex.

The effects of medicines such as ciclosporin, mycophenolate mofetil and tacrolimus (**medicines used to suppress the immune system**) may be reduced by Sevelamer carbonate Rowex. Your doctor will advise you if you are taking these medicines.

Thyroid hormone deficiency may uncommonly be observed in certain people taking **levothyroxine** (used to treat low thyroid hormone levels) and Sevelamer carbonate Rowex. Therefore your doctor may monitor the levels of thyroid stimulating hormone in your blood more closely.

If you are taking medicines to treat heartburn, gastroesophageal reflux disease (GERD) or gastric ulcers, such as **omeprazole, pantoprazole, or lansoprazole**, you should consult your doctor when taking Sevelamer carbonate Rowex.

Your doctor will check for interactions between Sevelamer carbonate Rowex and other medicines on a regular basis.

In some cases where Sevelamer carbonate Rowex should be taken at the same time as another medicine, your doctor may advise you to take this medicine 1 hour before or 3 hours after Sevelamer carbonate Rowex intake, or they may consider monitoring the blood levels of that medicine.

### Pregnancy, breast-feeding and fertility

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. It is unknown whether Sevelamer carbonate Rowex has any effect on unborn babies.

Tell your doctor if you wish to breast-feed your baby. It is unknown whether Sevelamer carbonate Rowex may pass through breast milk and affect your baby.

### Driving and using machines

Sevelamer carbonate Rowex is unlikely to affect your ability to drive or to use machines.

### Sevelamer carbonate Rowex contains lactose

Sevelamer carbonate Rowex contains **lactose** (milk sugar). If you have been told by your doctor that you have an **intolerance to some sugars**, contact your doctor before taking this medicinal product.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

### 3. How to take Sevelamer carbonate Rowex

Always take this medicine exactly as your doctor has told you. Check with your doctor if you are not sure. They will base the dose on your serum phosphorus level.

The recommended starting dose of Sevelamer carbonate Rowex tablets for adults and the elderly (> 65 years) is **one to two tablets of 800 mg with each meal, 3 times a day.**

**The tablets must be swallowed whole.** Do not crush, chew or break into pieces.

Initially your doctor will check the levels of phosphorus in your blood every 2-4 weeks and may adjust the dose of Sevelamer carbonate Rowex when necessary to reach an adequate phosphate level.

Patients taking Sevelamer carbonate Rowex should adhere to their prescribed diets.

#### **If you take more Sevelamer carbonate Rowex than you should**

In the event of a possible overdose you should contact your doctor immediately.

#### **If you forget to take Sevelamer carbonate Rowex**

If you have missed one dose, this dose should be omitted and the next dose should be taken at the usual time with a meal. Do not take a double dose to make up for a forgotten dose.

### 4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Since constipation may be an early symptom of a blockage in your intestine, please inform your doctor or pharmacist.

The following side effects have been reported in patients taking Sevelamer carbonate Rowex:

Very common (may affect more than 1 in 10 people):

Vomiting • constipation • upper abdominal pain • nausea

Common (may affect up to 1 in 10 people):

Diarrhoea • abdominal pain • indigestion • flatulence

Very rare (may affect up to 1 in 10000 people):

Hypersensitivity

Not known (frequency cannot be estimated from the available data):

Cases of itching, rash, slow intestine motility (movement)/blockages in the intestine, and perforation in the intestine wall have been reported.

#### **Reporting of side effects**

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via HPRA Pharmacovigilance, Earlsfort Terrace, IRL - Dublin 2; Tel: +353 1 6764971; Fax: +353 1 6762517.

Website: [www.hpra.ie](http://www.hpra.ie); E-mail: [medsafety@hpra.ie](mailto:medsafety@hpra.ie). By reporting side effects you can help provide more information on the safety of this medicine.

### 5. How to store Sevelamer carbonate Rowex

Keep this medicine out of the sight and reach of children.

*[For HDPE bottles without the carton]*

Do not use this medicine after the expiry date stated on the bottle after the letters "EXP". The expiry date refers to the last day of that month.

*[For HDPE bottles with the carton]*

Do not use this medicine after the expiry date stated on the bottle and carton after the letters "EXP". The expiry date refers to the last day of that month.

This medicinal product does not require any special storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

## 6. Contents of the pack and other information

#### **What Sevelamer carbonate Rowex contains**

- The active substance is sevelamer carbonate. Each film-coated tablet contains 800 mg of sevelamer carbonate.
- The other ingredients are lactose monohydrate, silica (colloidal anhydrous), zinc stearate. The tablet coating contains hypromellose (E464), diacetylated monoglycerides.

#### **What Sevelamer carbonate Rowex looks like and contents of the pack**

Sevelamer carbonate Rowex are oval, white to off-white film-coated tablets with the inscription 'SVL' on one side.

HDPE bottles with a polypropylene cap.

Each bottle contains 180, 200 or 210 tablets.

Packs containing 1, 2 or 3 bottles are available.

The HDPE bottles contain a desiccant. Do not remove this desiccant from the bottle.

Not all pack sizes may be marketed.

#### **Marketing Authorisation Holder and Manufacturers**

##### **Marketing Authorisation Holder**

Rowex Ltd., Bantry, Co. Cork, Ireland.

##### **Manufacturers**

Synthon Hispania S.L., Castello 1, Poligono Las Salinas, 08830 Sant Boi de Llobregat, Spain.

Salutas Pharma GmbH, Otto-von-Guericke-Allee 1, 39179 Barleben, Germany.

#### **This medicinal product is authorised in the Member States of the EEA under the following names:**

Denmark	Sevelamer Sandoz
Austria	Sevelamer Sandoz 800 mg Filmtabletten
Belgium	Sevelamer carbonate Sandoz 800 mg filmomhulde tabletten
Croatia	Sevelamer Sandoz 800 mg filmom obložene tablete
Cyprus	Sevelamer Carbonate Sandoz
Bulgaria	Sevelamer Sandoz
Czech Republic	Sevelamer Carbonate Sandoz 800 mg
Finland	Sevelamer Sandoz 800 mg tabletti, kalvopäällysteinen
France	SEVELAMER SANDOZ 800 mg, comprimé pelliculé
Ireland	Sevelamer carbonate Rowex 800 mg Film-coated tablets
Latvia	Sevelamer Carbonate Sandoz 800 mg Apvalkotās tableti
Lithuania	Sevelamer Sandoz 800 mg plėvele dengtos tabletės
Luxembourg	Sevelamer Sandoz 800 mg comprimés pelliculés
The Netherlands	Sevelameercarbonaat Sandoz 800 mg, filmomhulde tabletten
Norway	Sevelamer Sandoz 800 mg filmdrasjerte tablett
Poland	Sevelamer carbonate Sandoz
Romania	Carbonat de sevelamer Sandoz 800 mg comprimate filmate
Slovak Republic	Sevelamer carbonate Sandoz 800 mg
Slovenia	Sevelamer Lek 800 mg filmsko obložene tablete
Sweden	Sevelamer Sandoz 800 mg filmdragerade tablett

**This leaflet was last revised in 05/2018.**

I.M. L/950a 06-19