

## Ezetimibe Rowa 10mg Tablets

ezetimibe

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

### What is in this leaflet:

1. What Ezetimibe Rowa is and what it is used for
2. What you need to know before you take Ezetimibe Rowa
3. How to take Ezetimibe Rowa
4. Possible side effects
5. How to store Ezetimibe Rowa
6. Contents of the pack and other information

**ROWA**<sup>®</sup>  
Pharmaceuticals Limited

## 1 What Ezetimibe Rowa is and what it is used for

Ezetimibe Rowa is a medicine to lower increased levels of cholesterol.

Ezetimibe Rowa lowers levels of total cholesterol, “bad” cholesterol (LDL cholesterol), and fatty substances called triglycerides in the blood. In addition, Ezetimibe Rowa raises levels of “good” cholesterol (HDL cholesterol).

Ezetimibe, the active ingredient of Ezetimibe Rowa works by reducing the cholesterol absorbed in your digestive tract.

Ezetimibe Rowa adds to the cholesterol-lowering effect of statins, a group of medicines that reduce the cholesterol your body makes by itself.

Cholesterol is one of several fatty substances found in the bloodstream. Your total cholesterol is made up mainly of LDL and HDL cholesterol.

LDL cholesterol is often called “bad” cholesterol because it can build up in the walls of your arteries forming plaque. Eventually this plaque build-up can lead to a narrowing of the arteries. This narrowing can slow or block blood flow to vital organs such as the heart and brain. This blocking of blood flow can result in a heart attack or stroke.

HDL cholesterol is often called “good” cholesterol because it helps keep the bad cholesterol from building up in the arteries and protects against heart disease.

Triglycerides are another form of fat in your blood that may increase your risk for heart disease.

It is used for patients who cannot control their cholesterol levels by cholesterol lowering diet alone. You should stay on your cholesterol lowering diet while taking this medicine.

### Ezetimibe Rowa is used in addition to your cholesterol lowering diet if you have:

- a raised cholesterol level in your blood (primary hypercholesterolaemia [heterozygous familial and non-familial])
  - together with a statin, when your cholesterol level is not well controlled with a statin alone
  - alone, when statin treatment is inappropriate or is not tolerated
- a hereditary illness (homozygous familial hypercholesterolaemia) that increases the cholesterol level in your blood. You will also be prescribed a statin and may also receive other treatments.

If you have heart disease, Ezetimibe Rowa combined with cholesterol-lowering medicines called statins reduces the risk of heart attack, stroke, surgery to increase heart blood flow, or hospitalization for chest pain.

Ezetimibe Rowa does not help you lose weight.

## 2 What to know before you take Ezetimibe Rowa

If you use Ezetimibe Rowa together with a statin, please read the package leaflet of that particular medicine.

### Do not take Ezetimibe Rowa

- if you are allergic to ezetimibe or any of the other ingredients of this medicine (listed in section 6: Contents of the pack and other information).

### Do not take Ezetimibe Rowa together with a statin

- if you currently have liver problems
- if you are pregnant or breast-feeding.

### Warnings and precautions

Talk to your doctor or pharmacist before taking Ezetimibe Rowa

- Tell your doctor about all your medical conditions including allergies
- Your doctor should do a blood test before you start taking Ezetimibe Rowa with a statin. This is to check how well your liver is working.
- Your doctor may also want you to have blood tests to check how well your liver is working after you start taking Ezetimibe Rowa with a statin.

If you have moderate or severe liver problems, Ezetimibe Rowa is not recommended.

The safety and efficacy of the combined use of Ezetimibe Rowa and certain cholesterol lowering medicines, the fibrates have not been established.

### Children and adolescents

Do not give this medicine to children and

adolescents (6 to 17 years of age) unless prescribed by a specialist because there are limited data on safety and efficacy.

Do not give this medicine to children less than 6 years old because there is no information in this age group.

### Other medicines and Ezetimibe Rowa

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines. In particular, tell your doctor if you are taking medicine(s) with any of the following active ingredients:

- ciclosporin (often used in organ transplant patients)
- medicines with an active ingredient to prevent blood clots, such as warfarin, phenprocoumon, acenocoumarol or fluindione (anticoagulants)
- colestyramine (also used to lower cholesterol), because it affects the way Ezetimibe Rowa works
- fibrates (also used to lower cholesterol).

### Pregnancy and breast-feeding

Do not take Ezetimibe Rowa with a statin if you are pregnant, are trying to get pregnant or think you may be pregnant. If you get pregnant while taking Ezetimibe Rowa with a statin, stop taking both medicines immediately and tell your doctor.

There is no experience from the use of Ezetimibe Rowa without a statin during pregnancy. Ask your doctor for advice before using Ezetimibe Rowa if you are pregnant.

Do not take Ezetimibe Rowa with a statin if you are breast-feeding, because it is not known if the medicines are passed into breast milk.

Ezetimibe Rowa without a statin should not be used if you are breast-feeding. Ask your doctor for advice.

Ask your doctor or pharmacist for advice before taking any medicine.

### Driving and using machines

Ezetimibe Rowa is not expected to interfere with your ability to drive or to use machinery. However, it should be taken into account that some people may get dizzy after taking Ezetimibe Rowa.

### Ezetimibe Rowa contains lactose

Ezetimibe Rowa tablets contain a sugar called lactose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

### Ezetimibe Rowa contains sodium

This medicine contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially ‘sodium free’.

## 3 How to take Ezetimibe Rowa

Always take this medicine exactly as your doctor has told you. Continue taking your other cholesterol-lowering medicines unless your doctor tells you to stop. Check with your doctor or pharmacist if you are not sure.

- Before starting to take Ezetimibe Rowa, you should be on a diet to lower your cholesterol
- You should keep on this cholesterol lowering diet whilst taking Ezetimibe Rowa.

The recommended dose is one Ezetimibe Rowa 10 mg tablet, by mouth, once a day.

Take Ezetimibe Rowa at any time of the day. You can take it with or without food.

If your doctor has prescribed Ezetimibe Rowa along with a statin, both medicines can be taken at the same time. In this case, please read the dosage instructions in the package leaflet of that particular medicine.

If your doctor has prescribed Ezetimibe Rowa along with another medicine for lowering cholesterol containing the active ingredient colestyramine or any other medicine containing bile acid sequestrant, you should take Ezetimibe Rowa at least 2 hours before or 4 hours after taking the bile acid sequestrant.

### If you take more Ezetimibe Rowa than you should

Contact your doctor or pharmacist.

### If you forget to take Ezetimibe Rowa

Do not take a double dose to make up for a forgotten tablet, just take your normal amount of Ezetimibe Rowa at the usual time the next day.

### If you stop taking Ezetimibe Rowa

Talk to your doctor or pharmacist because your cholesterol may rise again.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

## 4 Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

The following terms are used to describe how often side effects have been reported:

- Very common (may affect more than 1 of 10 patients)
- Common (may affect up to 1 of 10 patients)
- Uncommon (may affect up to 1 of 100 patients)
- Rare (may affect up to 1 of 1,000 patients)
- Very rare (may affect up to 1 of 10,000 patients, including isolated reports).

**Contact your doctor immediately if you experience unexplained muscle pain, tenderness, or weakness. This is because on rare occasions, muscle problems, including muscle breakdown resulting in kidney damage, can be serious and may become a potentially life-threatening condition.**

Allergic reactions, including swelling of the face, lips, tongue, and/or throat that may cause difficulty in breathing or swallowing (which requires treatment right away) have been reported in general use.

When used alone, the following side effects were reported:

Common: abdominal pain; diarrhoea; flatulence; feeling tired

Uncommon: elevations in some laboratory blood tests of liver (transaminases) or muscle (CK) function; cough; indigestion; heartburn; nausea; joint pain; muscle spasms; neck pain; decreased appetite, pain, chest pain, hot flush; high blood pressure.

Additionally, when used with a statin, the following side effects were reported:

Common: elevations in some laboratory blood tests of liver function (transaminases); headache; muscle pain, tenderness or weakness

Uncommon: tingling sensation; dry mouth; itching, rash, hives; back pain; muscle weakness; pain in arms and legs; unusual tiredness or weakness; swelling, especially in the hands and feet.

When used with fenofibrate, the following side effect was reported: abdominal pain.

Additionally, the following side effects have been reported in general use: dizziness; muscle aches; liver problems; allergic reactions including rash and hives; raised red rash, sometimes with target-shaped lesions (erythema multiforme); muscle pain, tenderness or weakness; muscle breakdown; gallstones or inflammation of the gallbladder (which may cause abdominal pain, nausea, vomiting); inflammation of the pancreas often with severe abdominal pain; constipation, reduction in blood cell counts, which may cause bruising/bleeding (thrombocytopaenia), tingling sensation, depression: unusual tiredness or weakness; shortness of breath.

### Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via HPRA Pharmacovigilance.

Website: [www.hpra.ie](http://www.hpra.ie). By reporting side effects you can help provide more information on the safety of this medicine.

## 5 How to store Ezetimibe Rowa

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton or blister after "EXP." The expiry date refers to the last day of that month.

Do not store above 25°C. Store in the original package in order to protect from moisture.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

## 6 Contents of the pack and other information

### What Ezetimibe Rowa contains

- The active substance is ezetimibe. Each tablet contains 10 mg ezetimibe.
- The other ingredients are: lactose monohydrate, crospovidone (type B), copovidone K 28, sodium laurilsulfate, microcrystalline cellulose and magnesium stearate.

### What Ezetimibe Rowa looks like and contents of the pack

Ezetimibe Rowa tablets are white to off-white round convex tablets, of approximately 6 mm of diameter

#### Pack sizes:

Push-through blisters of transparent Triplex (PVC-PE-PVDC) /Aluminium in packs of 7, 10, 14, 20, 28, 30, 50, 84, 90, 98, 100, or 300 tablets.

Not all pack sizes may be marketed.

### Marketing Authorisation Holder and Manufacturer

Rowa Pharmaceuticals Ltd.

Bantry  
Co Cork  
Ireland

### Manufacturer

Laboratorios LICONSA, S.A.  
Polígono Industrial Miralcampo Avda.  
Miralcampo, N° 719200 Azuqueca de  
Henares  
Guadalajara  
SPAIN.

### This medicinal product is authorised in the Member States of the EEA under the following names:

Czech Republic	EZANTRIS
Finland	Ezetimib Liconsa 10 mg tabletti
France	Ézétimibe Almus 10 mg comprimé
Germany	Ezetimib AXiomed 10 mg tabletten
Ireland	Ezetimibe Rowa 10 mg Tablets
Poland	Ezetimibe Genoptim
Slovakia	EZENTRIS 10 mg
Spain	Ezetimiba Almus 10 mg comprimidos EFG
United Kingdom	Ezetimibe 10 mg tablets

This leaflet was last revised in 12/2020.

I.M. L/980a 12-20